



Review of the genus *Lasiochira* Meyrick, 1931 (Lepidoptera: Oecophoridae)

A. H. YIN¹, S. X. WANG^{1,3} & K. T. PARK²

¹College of Life Sciences, Nankai University, Tianjin 300071, China

²The Korean Academy of Science and Technology, Seongnam, Gyeonggi, 463-808 Korea. E-mail: ktpark02@gmail.com

³Corresponding author. E-mail: shxwang@nankai.edu.cn

Abstract

The genus *Lasiochira* Meyrick, 1931 is reviewed, based on the specimens collected from China, Korea, and Vietnam. Of the eight species involved in this study, six are described as new: *L. flavaterminata* sp. nov., *L. jianfengensis* sp. nov., *L. jiulongshana* sp. nov., *L. pallidiptera* sp. nov., *L. rosataenia* sp. nov. and *L. taiwanensis* sp. nov. Photographs of adults and genital structures as well as the wing venation are provided, along with a key to all the known species.

Key words: Lepidoptera, Oecophoridae, *Lasiochira*, taxonomy, new species

Introduction

Lasiochira Meyrick, 1931 is a small genus with two known species that are distributed in China. The type species, *L. camaropa* Meyrick, 1931, was described from Sichuan; *L. xanthacma* (Meyrick, 1938) was described from Yunnan and was also recorded to occur in Guangdong, Guizhou, Henan, and Shaanxi (Wang, 2006). Since Meyrick, no additional species have been added to this genus. We identified six new species of *Lasiochira* in this study, which were collected from China, Korea, and Vietnam.

Lasiochira is characterized by the forewing having a pale N-shaped pattern consisting of three joined stripes on the distal 3/5. It has the combination of the following characters: head with appressed scales, labial palpus with third segment slightly shorter than the second, antenna shorter than the forewing length and bearing pecten on scape; forewing with small tufts of erect scales, R_4 and R_5 stalked for about 2/3 length, M_1 to CuA_1 almost evenly spaced at base (Fig. 1); hindwing with anal area projected, R_s and M_1 separate and parallel, M_2 , M_3 and CuA_1 from the lower angle of the cell (Fig. 1); abdomen with five tergites densely spined in female (Fig. 2), six tergites densely spined in male (Fig. 3); male genitalia with the uncus relatively small, gnathos often hooked terminally, valva usually with dense fine hairs on distal half, sacculus with or without a terminal triangular process, aedeagus with cornuti of short stout spines or plates; female genitalia with apophyses posteriores longer than apophyses anteriores, ostium bursae situated on the anterior margin of the eighth sternite, ductus bursae rather short with sclerotization, and the corpus bursae with a triangular signum.

Little is known about the biology of *Lasiochira*.

Material and methods

This study is based on specimens deposited in the following institutions: Insect Collection, College of Life Sciences, Nankai University, Tianjin, China (NKUM); Department of Life Sciences, Incheon National University, Incheon, Korea (INU); and Korea National Arboretum, Seoul, Korea (KNA). Genitalia mounting methods follow Li (2002).

Etymology. The epithet of this new species is from Latin *flava*-, meaning yellow, and *terminatus*, meaning terminal, referring to the yellow apex in the forewing.

***Lasiochira camaropa* Meyrick, 1931**

(Figs. 12, 18)

Lasiochira camaropa Meyrick, 1931: 71.

Material examined. 1♂, Mt. Qingcheng, Sichuan Province, 14 Aug. 1990, other data unknown (NKUM); 1♂, Dahonghai, Chongqing, 1200 m, 13 Aug. 2012, coll. Xiaofei Yang and Tengteng Liu (NKUM).

Diagnosis. This species is characterized by the forewing with a bright ochreous yellow spot near the base of dorsum (Fig. 12), and the valva truncate at the apex in the male genitalia (Fig. 18).

Distribution. China (Sichuan).

***Lasiochira xanthacma* (Meyrick, 1938)**

(Figs. 13, 19, 26)

Allotalanta xanthacma Meyrick, 1938: 8.

Material examined. Henan Province: 7♂♂, 3♀♀, Neixiang County, 1350 m, 13–15 Jul. 1998, coll. Houhun Li; 6♂♂, 12♀♀, Mt. Baiyun, Song County, 1560 m, 22–24 Jul. 2001, coll. Dandan Zhang; 17♂♂, 10♀♀, Mt. Baiyun, Song County, 1580 m, 18–24 Jul. 2002, coll. Xinpu Wang; Shaanxi Province: 4♂♂, Feng County, 9–14 Jul. 1998, coll. Houhun Li; 1♂, Ankang City, 800 m, 4 Jul. 2003, coll. Haili Yu; Guizhou Province: 1♂, Mt. Fanjing, 1300 m, 3 Aug. 2001, coll. Houhun Li and Xinpu Wang; Guangdong Province: 2♀♀, Mt. Dadong, Lianzhou City, 650 m, 22, 24 Jun. 2004, coll. Dandan Zhang; Shanxi Province: 9♀♀, 1♂, Dahelinchang, 1340 m, 12–14 Jul. 2012, coll. Qiang Gao and Na Chen; 1♀, 3♂♂, Xiachuangdongxia, 1620 m, 18 Jul. 2012, coll. Qiang Gao and Na Chen; 1♀, 1♂, Xiachuanxixia, 1500 m, 16 Jul. 2012, coll. Qiang Gao and Na Chen; 1♀, Xiachuanbaohuzhan, 1600 m, 19 Jul. 2012, coll. Qiang Gao and Na Chen. (All deposited in NKUM).

Diagnosis. This species can be distinguished from its congeners by the sacculus without a distal process (Fig. 19). It is more similar to *L. pallidiptera* sp. nov. superficially (Fig. 13), but can be separated by the ovate ostium bursae open anteriorly (Fig. 26), whereas it is closed in the latter species.

Distribution. China (Guangdong, Guizhou, Henan, Shaanxi, Yunnan).

Acknowledgements

We express our cordial thanks to Prof. Y. S. Bae (Department of Life Science, Incheon National University, Incheon, Korea) and Dr. B. W. Lee (Korea National Arboretum, Pocheon, Korea) for the loan of specimens. This project is supported by the National Natural Science Foundation of China (No. 312111286 and No. 31372241), and in part by the National Institute of Biological Resources, Ministry of Environment, Korea (NIBR No. 2013-02-001).

References

- Meyrick, E. (1931) Second contribution to our knowledge about the Pyralidae and Microlepidoptera of Kwanhsien. *Bulletin de la Section Scientifique Académie Roumaine*, 14, 59–75.
- Meyrick, E. (1938) Materialien zu einer Mikrolepidopterenfauna des Yülingshanmassivs (Provinz Yünnan). *Deutsche Entomologische Zeitschrift*, 52, 1–29.
- Li, H.H. (2002) *The Gelechioidea of China (I) (Lepidoptera: Gelechioidea)*. Nankai University Press, Tianjin, 504 pp.
- Wang, S.X. (2006) *Oecophoridae of China (Insecta: Lepidoptera)*. Science Press, Beijing, 258 pp.